

兔抗 MNDA 多克隆抗体

中文名称：兔抗 MNDA 多克隆抗体

英文名称：Anti-MNDA rabbit polyclonal antibody

别名：PYHIN3

储存：冷冻（-20℃）避光

抗原：MNDA

宿主：Rabbit

反应种属：Human

相关类别：一抗

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	The myeloid cell nuclear differentiation antigen (MNDA) is detected only in nuclei of cells of the granulocyte-monocyte lineage. A 200-amino acid region of human MNDA is strikingly similar to a region in the proteins encoded by a family of interferon-inducible mouse genes, designated Ifi-201, Ifi-202, and Ifi-203, that are not regulated in a cell- or tissue-specific fashion. The 1.8-kb MNDA mRNA, which contains an interferon-stimulated response element in the 5-prime untranslated region, was significantly upregulated in human monocytes exposed to interferon alpha. MNDA is located within 2,200 kb of FCER1A, APCS, CRP, and SPTA1. In its pat
--------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	tern of expression and/or regulation, MNDA resembles IFI16, suggesting that these genes participate in blood cell-specific responses to interferons.
Applications:	WB
Name of antibody:	MNDA
Immunogen:	Synthesized peptide derived from C-terminal of human MNDA.
Full name:	myeloid cell nuclear differentiation antigen
Synonyms :	PYHIN3
SwissProt:	P41218
WB Predicted band size:	46 kDa
WB Positive control:	LOVO cells lysate
WB Recommended dilution:	500-3000

