

SMAD5 抗原(重组蛋白)

中文名称: SMAD5 抗原(重组蛋白)

- 英文名称: SMAD5 Antigen (Recombinant Protein)
- 储存: 冷冻(-20℃)
- 相关类别: 抗原

概述

Fusion protein corresponding to a region derived from C terminal 300 amino acids of human SMAD5

技术规格

Full name:	SMAD family member 5
Synonyms:	DWFC; JV5-1; MADH5
Swissprot:	Q99717
Gene Accession:	BC009682
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The protein encoded by this gene is involved in the TGF-beta sign aling pathway that results in an inhibition of the proliferation of he matopoietic progenitor cells. The encoded protein is activated by B MP type 1 receptor kinase. Three transcript variants encoding the s ame protein have been found for this gene. Transcriptional modula tor activated by BMP (bone morphogenetic proteins) type 1 recept or kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD). May for m trimers with the co-SMAD SMAD4. Interacts with PEBP2-alpha su



ł	ounit and	SMURF1.	Interacts	with	SUV39H1	and	SUV39H2.	Interacts
(via MH2	domain) v	with LEMD	93. In	teracts wi	th W	WP1.	