



兔抗 IFT57 多克隆抗体

中文名称: 兔抗 IFT57 多克隆抗体

英文名称: Anti-IFT57 rabbit polyclonal antibody

别名: intraflagellar transport 57; HIPPI; MHS4R2; ESRRBL1

储存: 冷冻 (-20°C)

抗原: IFT57

宿主: Rabbit

反应种属: Human, Mouse

相关类别 一抗

标记物 Unconjugate

克隆类型: rabbit polyclonal

技术规格

| | |
|----------------------|---|
| Background: | Required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro-apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'-AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear. |
| Applications: | ELISA, IHC |

| | |
|------------------------------------|-------------------------------|
| Name of antibody: | IFT57 |
| Immunogen: | Fusion protein of human IFT57 |
| Full name: | intraflagellar transport 57 |
| Synonyms: | HIPPI; MHS4R2; ESRRBL1 |
| SwissProt: | Q9NWB7 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human tonsil |
| IHC Recommend dilution: | 50-300 |

