

兔抗 IMPA1 多克隆抗体

中文名称：兔抗 IMPA1 多克隆抗体

英文名称：Anti-IMP1 rabbit polyclonal antibody

别名：IMP; IMPA

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：IMP1

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes an enzyme that dephosphorylates myo-inositol monophosphate to generate free myo-inositol, a precursor of phosphatidylinositol, and is therefore an important modulator of intracellular signal transduction via the production of the second messengers myo-inositol 1,4,5-trisphosphate and diacylglycerol. This enzyme can also use myo-inositol-1,3-diphosphate, myo-inositol-1,4-diphosphate, scyllo-inositol-phosphate, glucose-1-phosphate, glucose-6-phosphate, fructose-1-phosphate, beta-glycerophosphate, and 2'-AMP as substrates. This enzyme shows magnesium-dependent phosphatase activity and is inhibited by therapeutic concentrations of lithium. Inhibition of inositol monophosphate hydrolysis and s
--------------------	---

	ubsequent depletion of inositol for phosphatidylinositol synthesis may explain the anti-manic and anti-depressive effects of lithium administered to treat bipolar disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A pseudo gene of this gene is also present on chromosome 8q21.13.
Applications:	ELISA, WB, IHC
Name of antibody:	IMPA1
Immunogen:	Fusion protein of human IMPA1
Full name:	inositol monophosphatase 1
Synonyms:	IMP; IMPA
SwissProt:	P29218
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and Human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	30 kDa
WB Positive control:	Human cerebrum tissue lysate
WB Recommended dilution:	500-2000



