

兔抗 USO1 多克隆抗体

中文名称：兔抗 USO1 多克隆抗体

英文名称：Anti-USO1 rabbit polyclonal antibody

别名：TAP; VDP; P115

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：USO1

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is a peripheral membrane protein which recycles between the cytosol and the Golgi apparatus during interphase. It is regulated by phosphorylation: dephosphorylated protein associates with the Golgi membrane and dissociates from the membrane upon phosphorylation. Ras-associated protein 1 recruits this protein to coat protein complex II (COPII) vesicles during budding from the endoplasmic reticulum, where it interacts with a set of COPII vesicle-associated SNAREs to form a cis-SNARE complex that promotes targeting t

	o the Golgi apparatus. Alternative splicing results in multiple transcript variants.
Applications:	ELISA, WB, IHC
Name of antibody:	USO1
Immunogen:	Fusion protein of human USO1
Full name:	USO1 vesicle transport factor
Synonyms:	TAP; VDP; P115
SwissProt:	O60763
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	108 kDa
WB Positive control:	Mouse testis tissue , Jurkat and A549 cell lysates
WB Recommended dilution:	500-2000





