

## 兔抗 PAPPA-AS1 多克隆抗体

中文名称: 兔抗 PAPPA-AS1 多克隆抗体

英文名称: Anti-PAPPA-AS1 rabbit polyclonal antibody

别 名: DIPAS; TAF4B; PAPPAS; TAF2C2; PAPPAAS; PAPPA-AS; TAFII105; NCRNA00156

相关类别: 一抗

抗 原: PAPPA-AS1

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

PAPP A was originally isolated from pregnancy serum as heterotetrameric protein complex with molecular weight a pproximately 500 kDa, consisting of two PAPP A subunits disulfide-linked with two subunits of the proform of eosi nophil major basic protein (proMBP). In such a complex proMBP functions as an inhibitor of PAPP A proteinase a ctivity. PAPP A is widely used as a marker of some path ologies during pregnancy. Reduced level of PAPP A in pregnancy blood in the first trimester is associated with fet al Down syndrome. PAPP A is considered as one of the best biochemical markers in early pregnancy used to scre



|                                    | en for Down syndrome during the first trimester (Wald, N. J. et al).  |
|------------------------------------|---|
| Applications:                      | ELISA, IHC  |
| Name of antibody:                  | PAPPA-AS1   |
| Immunogen:                         | Synthetic peptide of human PAPPA-AS1                                  |
| Full name:                         | PAPPA antisense RNA 1   |
| Synonyms:                          | DIPAS; TAF4B; PAPPAS; TAF2C2; PAPPAAS; PAPPA-AS; TAFII105; NCRNA00156 |
| SwissProt:                         | Q5QFB9  |
| <b>ELISA Recommended dilution:</b> | 5000-10000  |
| IHC positive control:              | Human tonsil  |
| IHC Recommend dilution:            | 20-100  |

