

兔抗 ITGB4(Ab-1510) 多克隆抗体

中文名称: 兔抗 ITGB4(Ab-1510) 多克隆抗体

英文名称: Anti-ITGB4(Ab-1510) rabbit polyclonal antibody

别 名: CD104

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: ITGB4(Ab-1510)

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

| Name of antibody: | ITGB4(Ab-1510) |
|-------------------|---|
| Applications: | WB, IHC |
| | bunit, a receptor for the laminins. |
| Background: | n and cell growth. This gene encodes the integrin beta 4 su |
| | hesion, and transduced signals that regulate gene expressio |
| | ding specificities. Integrins mediate cell-matrix or cell-cell ad |
| | a polypeptides form complexes that vary in their ligand-bin |
| | oprotein receptors. Different combinations of alpha and bet |
| | units, that are noncovalently associated transmembrane glyc |
| | Integrins are heterodimers comprised of alpha and beta sub |



| Immunogen: | Synthesized non-phosphopeptide derived from human ITGB4 around the phosphorylation site of tyrosine 1510 (R-D-Y(p)-S-T). |
|--------------------------|--|
| Full name: | integrin, beta 4 |
| Synonyms : | CD104 |
| SwissProt: | P16144 |
| IHC positive control: | Human breast carcinoma tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 202 kDa |
| WB Positive control: | HepG2 cells treated with PMA (125ng/ml, 30mins) lysate |
| WB Recommended dilution: | 500-3000 |





