

## 兔抗 PDIA4 多克隆抗体

中文名称: 兔抗 PDIA4 多克隆抗体

英文名称: Anti-PDIA4 rabbit polyclonal antibody

别 名: protein disulfide isomerase family A member 4; ERP70; ERP72; ERp-72

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: PDIA4

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

Background:	This gene encodes a member of the disulfide isomerase (P DI) family of endoplasmic reticulum (ER) proteins that cata lyze protein folding and thiol-disulfide interchange reactio ns. The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, t wo TRX-like domains and a C-terminal ER-retention sequence. This protein, when bound to cyclophilin B, enhances t he rate of immunoglobulin G intermolecular disulfide bond
	ing and antibody assembly.
Applications:	ELISA, WB, IHC
Name of antibody:	PDIA4



Immunogen:	Fusion protein of human PDIA4
Full name:	protein disulfide isomerase family A member 4
Synonyms:	ERP70; ERP72; ERp-72
SwissProt:	P13667
ELISA Recommended dilution :	5000-10000
IHC positive control:	Human cervical cancer and Human thyroid cancer
IHC Recommend dilution:	30-150
WB Predicted band size:	73 kDa
WB Positive control:	NIH/3T3 and RAW264.7 cell lysates
WB Recommended dilution:	500-2000







