

兔抗 SLC3A2 多克隆抗体

中文名称：兔抗 SLC3A2 多克隆抗体

英文名称：Anti-SLC3A2 rabbit polyclonal antibody

别名：solute carrier family 3 member 2; 4F2; CD98; MDU1; 4F2HC; 4T2HC; NACAE; CD98HC

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC3A2

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene is a member of the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through di-sulfide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intracellular calcium levels and transports L-type amino acids. Alternatively spliced transcript variants, encoding different isoforms, have been characterized.
Applications:	ELISA, WB, IHC
Name of antibody:	SLC3A2

Immunogen:	Fusion protein of human SLC3A2
Full name:	solute carrier family 3 member 2
Synonyms:	4F2; CD98; MDU1; 4F2HC; 4T2HC; NACAE; CD98HC
SwissProt:	P08195
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer and Human liver cancer
IHC Recommend dilution:	30-150
WB Predicted band size:	68 kDa
WB Positive control:	A375 and Jurkat cell lysates
WB Recommended dilution:	500-2000



