

兔抗 SLC7A5 多克隆抗体

- 中文名称: 兔抗 SLC7A5 多克隆抗体
- 英文名称: Anti-SLC7A5 rabbit ployclonal antibody
- 别 名: E16; CD98; LAT1; 4F2LC; MPE16; hLAT1; D16S469E
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: SLC7A5
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

a. Applications: ELISA, WB	ner ote LA und: er pla AT edi	ute carrier family 7 member 5 also known as large utral amino acids transporter small subunit 1 is a pr in that in humans is encoded by the SLC7A5gene. If is expressed predominantly in adult lung and liv but is also expressed in brain, thymus, retina, testis, centa, bone marrow and fetal liver. In the retina, L 1 localizes to the blood-retinal-barrier (BRB) and m ates L-leucine transport from the blood to the retin
	a.	•
Name of antibody: SLC7A5	-	



Immunogen:	Synthetic peptide of human SLC7A5
Full name:	solute carrier family 7 (amino acid transporter light ch ain, L system), member 5
Synonyms :	E16; CD98; LAT1; 4F2LC; MPE16; hLAT1; D16S469E
SwissProt:	Q01650
ELISA Recommended dilution:	1000-2000
WB Predicted band size:	55 kDa
WB Positive control:	823 cells
WB Recommended dilution:	200-1000

