

## 小鼠抗 PSMC3 单克隆抗体

中文名称: 小鼠抗 PSMC3 单克隆抗体

英文名称: Anti-PSMC3 mouse monoclonal antibody

别 名: proteasome 26S subunit, ATPase 3; TBP1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Mouse

抗 原: PSMC3

反应种属: Human

标 记 物: Unconjugate

克隆类型: mouse monoclonal

## 技术规格

**Background:** 

The 26S proteasome is a multicatalytic proteinase complex with a hi ghly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-ident ical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteaso mes are distributed throughout eukaryotic cells at a high concentrati on and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteas ome, the immunoproteasome, is the processing of class I MHC pept



	ides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. A pseudogene has been identified on chromosome 9.
Applications:	WB
Name of antibody:	PSMC3
Immunogen:	Fusion protein of human PSMC3
Full name:	proteasome 26S subunit, ATPase 3
Synonyms:	TBP1
SwissProt:	P17980
WB Predicted band size:	49 KD
WB Positive control :	293T, Hela, HepG2, A431 cell lysates
WB Recommended dilution:	200-1000