

## 兔抗 HEXB 多克隆抗体

中文名称：兔抗 HEXB 多克隆抗体

英文名称：Anti-HEXB rabbit polyclonal antibody

别 名：ENC-1AS; HEL-248

相关类别：一抗

抗 原：HEXB

储 存：冷冻（-20℃）避光

宿 主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：Unconjugate

### 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | Hexosaminidase B is the beta subunit of the lysosomal enzyme beta-hexosaminidase that, together with the cofactor GM2 activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. Beta-hexosaminidase is composed of two subunits, alpha and beta, which are encoded by separate genes. Both beta-hexosaminidase alpha and beta subunits are members of family 20 of glycosyl hydrolases. |
| <b>Applications:</b>     | WB, IHC  |
| <b>Name of antibody:</b> | HEXB   |
| <b>Immunogen:</b>        | Synthesized peptide derived from C-terminal of human HEXB.   |
| <b>Full name:</b>        | hexosaminidase B (beta polypeptide)  |
| <b>Synonyms :</b>        | ENC-1AS; HEL-248   |

|                                  |                              |
|----------------------------------|------------------------------|
| <b>SwissProt:</b>                | P07686                       |
| <b>IHC positive control</b><br>: | Human colon carcinoma tissue |
| <b>IHC Recommend dilution:</b>   | 50-100                       |
| <b>WB Predicted band size:</b>   | 63 kDa                       |
| <b>WB Positive control</b><br>:  | Jurkat cells lysate          |
| <b>WB Recommended dilution:</b>  | 500-3000                     |



