

兔抗 PTPRG 多克隆抗体

- 中文名称: 兔抗 PTPRG 多克隆抗体
- 英文名称: Anti-PTPRG rabbit polyclonal antibody
- 别名: PTPG; HPTPG; RPTPG; R-PTP-GAMMA
- 相关类别: 一抗
- 储存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗原: PTPRG
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are kno wn to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP po ssesses an extracellular region, a single transmembrane r egion, and two tandem intracytoplasmic catalytic domai ns, and thus represents a receptor-type PTP. The extrace llular region of this PTP contains a carbonic anhydrase-li ke (CAH) domain, which is also found in the extracellula r region of PTPRBETA/ZETA. This gene is located in a ch romosomal region that is frequently deleted in renal cell
-------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



	carcinoma and lung carcinoma, thus is thought to be a candidate tumor suppressor gene.
Applications:	ELISA, IHC
Name of antibody:	PTPRG
Immunogen:	Synthetic peptide of human PTPRG
Full name:	protein tyrosine phosphatase, receptor type, G
Synonyms:	PTPG; HPTPG; RPTPG; R-PTP-GAMMA
SwissProt:	P23470
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil and human cervical cancer
IHC Recommend dilution:	25-100

