

兔抗 RRAGA 多克隆抗体

中文名称：兔抗 RRAGA 多克隆抗体

英文名称：Anti-RRAGA rabbit ployclonal antibody

别 名：FIP1; RAGA; FIP-1

相关类别：一抗

储 存：冷冻（-20℃）

宿 主：Rabbit

抗 原：RRAGA

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | Ras-related GTP-binding protein A is a protein that in humans is encoded by the RRAGAgene. Guanine nucleotide-binding protein forming heterodimeric Rag complexes required for the amino acid-induced relocation of mTORC1 to the lysosomes and its subsequent activation by the GTPase RHEB. This is a crucial step in the activation of the TOR signaling cascade by amino acids. Involved in the RCC1/Ran-GTPase pathway. May play a direct role in a TNF-alpha signaling pathway leading to induction of cell death. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | RRAGA |

| | |
|------------------------------------|-------------------------------|
| Immunogen: | Fusion protein of human RRAGA |
| Full name: | Ras-related GTP binding A |
| Synonyms : | FIP1; RAGA; FIP-1 |
| SwissProt: | Q7L523 |
| ELISA Recommended dilution: | 2000-5000 |
| IHC positive control: | Human breast cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 37 kDa |
| WB Positive control: | Hepg2 and hela cells |
| WB Recommended dilution: | 500-2000 |



