

兔抗 MORC2 多克隆抗体

中文名称: 兔抗 MORC2 多克隆抗体

英文名称: Anti-MORC2 rabbit polyclonal antibody

别名: ZCW3; ZCWCC1

抗原: MORC2

储存: 冷冻 (-20℃)

宿主: Rabbit

反应种属: Human

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

The CW domain is a structural module found in many vertebrate, parasitic and plant proteins. It consists of a mononuclear four-cysteine zinc-finger domain that plays a role in DNA binding, chromatin methylation and early embryonic development. ZCWCC1 (zinc finger CW-type coiled-coil domain protein 1), also known as MORC2 (MORC family CW-type zinc finger protein 2) or ZCW3, is a 1,032 amino acid protein that contains one CW-type zinc finger domain. ZCWCC1 is located on chromosome 22 and is ubiquitously expressed with highest expression in pancreas, smooth muscle and testis. Expression of ZCWCC1 is upregulated in hypoxia, a pathological condition characterized by an in

| | |
|------------------------------------|---|
| | adequate supply of oxygen in the blood. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | MORC2 |
| Immunogen: | Fusion protein of human MORC2 |
| Full name: | MORC family CW-type zinc finger 2 |
| Synonyms: | ZCW3; ZCWCC1 |
| SwissProt: | Q9Y6X9 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human lung cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 118 kDa |
| WB Positive control: | 231, 293T, Hela and A172 cell lysates |
| WB Recommended dilution: | 500-2000 |



