

## 兔抗 ABCA1 多克隆抗体

中文名称：兔抗 ABCA1 多克隆抗体

英文名称：Anti-ABCA1 rabbit polyclonal antibody

别名：ATP binding cassette subfamily A member 1; TGD; ABC1; CERP; ABC-1; HDLDT1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：ABCA1

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

<b>Background:</b>	The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ABC1 subfamily. Members of the ABC1 subfamily comprise the only major ABC subfamily found exclusively in multicellular eukaryotes. With cholesterol as its substrate, this protein functions as a cholesterol efflux pump in the cellular lipid removal pathway. Mutations in this gene have been associated with Tang
--------------------	---

	ier's disease and familial high-density lipoprotein deficiency.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	ABCA1
<b>Immunogen:</b>	Synthetic peptide of human ABCA1
<b>Full name:</b>	ATP binding cassette subfamily A member 1
<b>Synonyms:</b>	TGD; ABC1; CERP; ABC-1; HDLDT1
<b>SwissProt:</b>	O95477
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human esophagus cancer
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	254 kDa
<b>WB Positive control:</b>	Human bladder transitional cell carcinoma grade 2-3 tissue, Human bladder transitional cell carcinoma cancer next level 2-3 tissue lysates
<b>WB Recommended dilution:</b>	200-1000



